COVID-19 VACCINE Priority Recommendations – March 19, 2021

Categories are broadly inclusive, but people who work directly with the public or in situations in which they cannot maintain masks and a distance of 6 feet from others should be prioritized. All Philadelphia vaccine sites are only for Philadelphia residents only.

Phase 1a

| HIGH RISK FOR EXPOSURE AND TRANSMISSION TO VULNERABLE PERSONS (PATIENT-FACING STAFF) | HIGH RISK OF MORBIDITY/MORTALITY |
|--|-----------------------------------|
| Hospital staff | Long-term care facility residents |
| COVID testing staff | |
| COVID vaccination & lab staff | |
| Long-term care facility staff | |
| Emergency medical services | |
| Home health care | |
| Prison health services | |
| Outpatient clinics, FQHCs | |
| Other healthcare providers | |

Phase 1b

| HIGH RISK FOR EXPOSURE AND PERFORM ESSENTIAL DUTIES | HIGH RISK OF MORBIDITY/MORTALITY |
|---|---|
| Police, paramedics/EMTs and firefighters | Persons living or working in high-risk congregate residential facilities: prisons, shelters for the homeless, residential drug and alcohol treatment, psychiatric facilities, rehabilitation, and specialized services housing (does not include apartments, condos, and other independent living facilities) |
| Service providers required to do home visits or visits to high-risk congregate settings, staff at senior centers/day programs for seniors and those with intellectual disabilities | Persons age 65+ |
| Public transit | Persons with the following high risk health |
| Food distribution, prep, or service workers Frontline child or public facing staff of childcare, K-12 schools, libraries and recreation centers. Includes custodians and food preparation staff. Public facing staff of high-volume retail settings Factory/manufacturing workers whose work conditions require them to be present in person at sites with 20 or more people per room and/or | conditions: Cancer Chronic kidney disease; Chronic obstructive pulmonary disease (COPD) Heart conditions, such as heart failure, coronary artery disease, or cardiomyopathies |

| those not able to consistently maintain 6 feet of | Immunocompromised state from solid |
|---|--|
| distance and masking while at work | organ or bone marrow transplant, immune |
| Clergy | deficiencies, HIV, use of oral |
| | corticosteroids, or use of other immune |
| | weakening medicines |
| | Obesity (body mass index [BMI] ≥30 |
| | Sickle cell disease |
| | Smoking |
| | Type 1 and 2 diabetes mellitus |
| | Down Syndrome |
| | Pregnancy |
| | People with intellectual disabilities |

Phase 1c

| OTHER ESSENTIAL WORKERS BUT WITH LOWER RISK OF EXPOSURE | HIGH RISK OF MORBIDITY/MORTALITY |
|--|----------------------------------|
| Sanitation workers | |
| Maintenance/janitorial workers (note that those working in schools and daycares are eligible in Phase 1B) | |
| Utility workers (except those frontline workers included as service providers who must do home visits in Phase 1B) | |
| Postal and package delivery workers | |
| Higher education | |
| Finance: public facing, non-remote positions only | |
| Transportation: airport and train workers, Taxi/rideshare drivers | |
| Construction | |
| IT & Telecommunications, Press | |
| Legal | |
| Public Health | |

Phase 2

Everyone \geq 16 years of age not previously immunized